

$SU(3)$ and CP violating weak and strong final state phases for B_d^0 and B_s^0 decays

Fayyazuddin

National Centre for Physics and Department of Physics
Quaid-i-Azam University
Islamabad, Pakistan.

February 2, 2008

Abstract

Using rotation in $SU(3)$ space, a set of relations between various decay modes of B_d and B_s are derived. The decays $\bar{B}_d^0 \rightarrow K^{*-}\pi^+(\rho^+K^-)$, $\bar{B}_s^0 \rightarrow K^{*-}K^+$ are expressed in terms of decay parameters of $\bar{B}_d^0 \rightarrow \rho^-\pi^+(\rho^+\pi^-)$. In particular the parameters $r_{-+}(r_{+-})$ of $B_d \rightarrow \rho\pi$ decays are obtained in terms of experimentally known decay rates $R_{-+}(R_{+-}) = \frac{1}{2}(\Gamma_{\rho^+\pi^-(\rho^-\pi^+)} + \bar{\Gamma}_{\rho^-\pi^+(\rho^+\pi^-)})$, $R'_{-+}(R'_{+-}) = \frac{1}{2}(\Gamma_{K^{*+}\pi^-(\rho^-K^+)} + \bar{\Gamma}_{K^{*-}\pi^+(\rho^+K^-)})$, known parameters $\bar{\lambda}$, $f_{K^*}/f_\rho(f_K/f_\pi)$ and two parameters $B_{-+} = \frac{R_{-+}}{|T^{-+}|^2}$, $B_{+-} = \frac{R_{+-}}{|T^{+-}|^2}$ which are determined by using factorization for tree amplitudes T^{-+} and T^{+-} . We find $r_{-+} = 0.21 \pm 0.04$, $r_{+-} = 0.25 \pm 0.06$. With these values the following bounds on ($z \equiv \cos \gamma \cos \delta$, $x = \sin \gamma \sin \delta$) are derived:

$$-0.34(-0.33) \leq z_{-+}(z_{+-}) \leq 0.28(0.27)$$

and

$$0.16(0.43) \leq x_{-+}(x_{+-}) \leq 0.58(1.00)$$

From (x, z) plot we obtain following bounds on weak phase γ and strong phases δ 's $z_{-+} > 0, \gamma \geq 70^\circ, 10^\circ \leq \delta_{-+} \leq 40^\circ, z_{-+} < 0, \gamma \geq 65^\circ, (180 - \delta_{-+})$. For δ_{+-} we get $25^\circ \leq \delta_{+-} \leq 90^\circ$ or $(180 - \delta_{+-})$.

The decays of B mesons to two light mesons which belong to an octet or nonet representation of $SU(3)$ have been extensively studied [1]. The various decay modes can be related to each other by rotation in $SU(3)$ space, using the relations [2]

$$[F_j^i, q_k] = \delta_k^i q_j - \frac{1}{3} \delta_j^i q_k \quad (1)$$

$$[F_j^i, q^k] = -\delta_j^k q^i + \frac{1}{3} \delta_j^i q^k \quad (2)$$

where F_j^i are $SU(3)$ generators. In particular, using

$$[F_2^3, H_W(s)] = H_W(d) \quad (3)$$

$$F_3^2 |\bar{s}\rangle = -|\bar{d}\rangle, \quad F_3^2 |d\rangle = |s\rangle$$

$$F_2^3 |\bar{d}\rangle = -|\bar{s}\rangle \quad (4)$$

$$F_2^3 |\bar{B}_d^0\rangle = F_2^3 |b\bar{d}\rangle = -|b\bar{s}\rangle = -|\bar{B}_s^0\rangle \quad (5)$$

we can relate various decay modes. The $SU(3)$ relations for various decay modes have been studied in references [3, 4]. In this paper, we concentrate on the decay modes which we believe have not been covered although there is bound to be overlap. Suppressing the CKM factors (which can be inserted for each relevant amplitude) we obtain the following relations for various decay

modes for $B \rightarrow PV$ decays:

$$\begin{aligned}
\langle K^{*0} \bar{K}^0 | H_W(d) | \bar{B}_d^0 \rangle &= \langle K^{*0} \bar{K}^0 | H_W(s) | \bar{B}_s^0 \rangle \\
&+ \left\{ \frac{1}{\sqrt{2}} [\langle \rho^0 \bar{K}^0 | H_W(s) | \bar{B}_d^0 \rangle - \langle \omega \bar{K}^0 | H_W(s) | \bar{B}_d^0 \rangle] \right. \\
&\quad \left. - \langle \phi \bar{K}^0 | H_W(s) | \bar{B}_d^0 \rangle \right\} \\
&= \frac{1}{2} [(C' - P' - P'_{EW}) - (C' + P' + \frac{1}{3} P'_{EW})] \\
&\quad - (-P' + \frac{1}{3} P'_{EW}) \}
\end{aligned} \tag{6}$$

$$\begin{aligned}
\langle K^0 \bar{K}^{*0} | H_W(d) | \bar{B}_d^0 \rangle &= + \left\{ \frac{1}{\sqrt{2}} \left[\begin{aligned} &\langle K^0 \bar{K}^{*0} | H_W(s) | \bar{B}_d^0 \rangle \\ &- \left[\begin{aligned} &\langle \pi^0 \bar{K}^{*0} | H_W(s) | \bar{B}_d^0 \rangle \\ &+ \langle \eta_{ns} \bar{K}^{*0} | H_W(s) | \bar{B}_d^0 \rangle \end{aligned} \right] \end{aligned} \right] \right\}
\end{aligned} \tag{7}$$

$$\text{where } \eta_{ns} = \frac{1}{\sqrt{3}} (\eta_8 + \sqrt{2} \eta_1), \quad \eta_s = \frac{1}{\sqrt{3}} (\eta_1 - \sqrt{2} \eta_8) \tag{8}$$

$$\langle K^{*+} K^- | H_W(d) | \bar{B}_d^0 \rangle = \langle K^{*+} K^- | H_W(s) | \bar{B}_s^0 \rangle - \langle \rho^+ K^- | H_W(s) | \bar{B}_d^0 \rangle \tag{9}$$

$$\langle K^{*-} K^+ | H_W(d) | \bar{B}_d^0 \rangle = \langle K^{*-} K^+ | H_W(s) | \bar{B}_s^0 \rangle - \langle \pi^+ K^{*-} | H_W(s) | \bar{B}_d^0 \rangle \tag{10}$$

$$\begin{aligned}
\langle K^{*0} K^- | H_W(d) | B^- \rangle &= \frac{1}{\sqrt{2}} [\langle \rho^0 K^- | H_W(s) | B^- \rangle - \langle \omega K^- | H_W(s) | B^- \rangle] \\
&\quad - \langle \phi K^- | H_W(s) | B^- \rangle \\
&= \frac{1}{2} \left[(T' + C' + P' + P'_{EW}) - \left(T' + C' + P' + \frac{1}{3} P'_{EW} \right) \right] \\
&\quad - \left[-P' + \frac{1}{3} P'_{EW} \right]
\end{aligned} \tag{11}$$

$$\begin{aligned}
\langle K^0 K^{*-} | H_W(d) | B^- \rangle &= \frac{1}{\sqrt{2}} [\langle \pi^0 K^{*-} | H_W(s) | B^- \rangle - \langle \eta_{ns} K^{*-} | H_W(s) | B^- \rangle] \\
&\quad + \langle \eta_s K^{*-} | H_W(s) | B^- \rangle
\end{aligned} \tag{12}$$

For the sake of completeness, using the above technique we derive the fol-

lowing relations .

$$\langle \rho^- \pi^+ | H_W(d) | \bar{B}_d^0 \rangle = \langle K^{*-} \pi^+ | H_W(s) | \bar{B}_d^0 \rangle + \langle \rho^- \pi^+ | H_W(s) | \bar{B}_s^0 \rangle \quad (13)$$

$$\langle \rho^+ \pi^- | H_W(d) | \bar{B}_d^0 \rangle = \langle \rho^+ K^- | H_W(s) | \bar{B}_d^0 \rangle + \langle \rho^+ \pi^- | H_W(s) | \bar{B}_s^0 \rangle \quad (14)$$

$$\langle \phi \pi^- | H_W(d) | B^- \rangle = \langle \phi K^- | H_W(s) | B^- \rangle + \langle \bar{K}^{*0} \pi^- | H_W(s) | B^- \rangle \quad (15)$$

$$\begin{aligned} \langle \phi \pi^0 | H_W(d) | \bar{B}_d^0 \rangle &= -\frac{1}{\sqrt{2}} \left(\langle \phi \bar{K}^0 | H_W(s) | \bar{B}_d^0 \rangle + \langle \bar{K}^{*0} \pi^0 | H_W(s) | \bar{B}_d^0 \rangle \right. \\ &\quad \left. + \langle \phi \pi^0 | H_W(s) | \bar{B}_d^0 \rangle \right) \\ &= -\frac{1}{\sqrt{2}} \left[-P' - \frac{1}{3} P'_{EW} \right] + \frac{1}{\sqrt{2}} [C' - P' + P'_{EW}] \\ &\quad + \frac{1}{\sqrt{2}} [-C' - P'_{EW}] \end{aligned} \quad (16)$$

We note that only penguin contributes to the decay channel $\bar{B}^0 \rightarrow K^{*0} \bar{K}^0$, $\bar{B}^0 \rightarrow K^0 \bar{K}^{*0}$. Hence it follows from Eqs.(6), (7), (11) and (12)

$$\begin{aligned} \langle K^{*0} \bar{K}^0 | H_W(d) | \bar{B}_d^0 \rangle &= P = |V_{ub}| |V_{ud}^*| e^{-i\gamma} P_u - |V_{cb}| |V_{cd}^*| P_c \\ &= |V_{cb}| |V_{cs}| |P_c| \times [-\bar{\lambda} + r e^{i(-\gamma+\delta)}] \end{aligned} \quad (17)$$

$$\begin{aligned} \langle K^{*0} \bar{K}^0 | H_W(s) | \bar{B}_s^0 \rangle &= P' = |V_{ub}| |V_{us}^*| e^{-i\gamma} P_u + |V_{cb}| |V_{cs}^*| P_c \\ &\approx |V_{cb}| |V_{cs}| |P_c| \end{aligned} \quad (18)$$

$$\bar{A}(\bar{B}_d^0 \rightarrow K^0 \bar{K}^{*0}) = |V_{cb}| |V_{cs}| \left| \tilde{P}_c \right| \times [-\bar{\lambda} + \tilde{r} e^{i(-\gamma+\delta)}] \quad (19)$$

$$\bar{A}(\bar{B}_s^0 \rightarrow K^0 \bar{K}^{*0}) \approx |V_{cb}| |V_{cs}| \left| \tilde{P}_c \right| \quad (20)$$

$$\bar{A}(B^- \rightarrow K^{*0} K^-) = |V_{cb}| |V_{cs}| |P_c^{0-}| \left[-\bar{\lambda} + r'_{0-} e^{i(-\gamma+\delta'_{0-})} \right] \quad (21)$$

$$\bar{A}(B^- \rightarrow \phi K^-) = |V_{cb}| |V_{cs}| |P_c^{0-}| \left[1 + \frac{1}{3} \delta_{EW}^{0-} \right] \quad (22)$$

$$\bar{A}(B^- \rightarrow K^0 K^{*-}) = |V_{cb}| |V_{cs}| |P_c'^{0-}| \left[-\bar{\lambda} + r'_{-0} e^{i(-\gamma+\delta'_{-0})} \right] \quad (23)$$

$$\bar{A}(B^- \rightarrow \eta_s K^{*-}) = |V_{cb}| |V_{cs}| |P_c'^{0-}| \left[1 + \frac{1}{3} \delta_{EW}^{0-} \right] \quad (24)$$

where

$$r e^{i\delta} = \frac{|V_{ub}| |V_{ud}^*|}{|V_{cb}| |V_{cs}|} \left| \frac{P_u}{P_c} \right| e^{i\delta} \quad (25)$$

$$\begin{aligned}
V_{cs} &= 1 - \frac{\lambda^2}{2}, V_{cd} = -\lambda, V_{us} = \lambda, V_{ud} = 1 - \frac{\lambda^2}{2} \\
\bar{\lambda} &= \frac{\lambda}{1 - \lambda^2/2}
\end{aligned} \tag{26}$$

and

$$\delta_{EW} = \frac{|P'_{EW}|}{|P'_c|}$$

It may also be noted, that for the decay channel $B^- \rightarrow \phi \pi^-$, only color suppressed and electroweak penguin contributes. Negelecting the color suppressed penguin contribution in Eq.(15), we get

$$\bar{A}(B^- \rightarrow \bar{K}^{*0} \pi^-) = |V_{cb}| |V_{cs}| |P_c'^{0-}| \tag{27}$$

and

$$\bar{A}(B^- \rightarrow \rho^- \bar{K}^0) = |V_{cb}| |V_{cs}| |P_c'^{-0}| \tag{28}$$

Electroweak penguins are in a class different from those of gluon penguins. Assumung factorization for electroweak penguin, we get from Eq.(6), an in-teresting sum rule

$$f_\rho F_1^{B-K}(m_\rho^2) - \frac{1}{3} f_\omega F_1^{B-K}(m_\omega^2) - \frac{2}{3} f_\phi F_1^{B-K}(m_\phi^2) = 0 \tag{29}$$

Assuming

$$F_1^{B-K}(m_\rho^2) = F_1^{B-K}(m_\omega^2) = F_1^{B-K}(m_\phi^2) \tag{30}$$

we obtain the relation

$$f_\rho - \frac{1}{3} f_\omega - \frac{2}{3} f_\phi = 0 \tag{31}$$

reminiscent of current algebra, and spectral function sum rules of 1960's [5]. Sum rule (31) is very well satisfied by the experimental values [6]

$$f_\rho = (209 \pm 1) MeV, f_\omega = 187 \pm 3 MeV, f_\phi = 221 \pm 3 MeV$$

The assumption stated in Eq (30) holds as the form factor F_1 at 1GeV will not differ much at the masses m_ρ^2, m_ω^2 and m_ϕ^2

Similarly we get the sum rule

$$f_\pi - \frac{1}{3} f_{\eta_{ns}} - \frac{2}{3} f_{\eta_s} = 0 \tag{32}$$

Finally, from the term in curly bracket in Eq.(6) we get the following relations between average decay rates and mixing induced CP asymmetries for $B_d^0 \rightarrow \rho^0 \bar{K}^0, \omega \bar{K}^0, \phi \bar{K}^0$

$$\frac{\langle \Gamma \rangle_{\omega K} + \langle \Gamma \rangle_{\rho^0 K}}{\langle \Gamma \rangle_{\phi K}} \approx 1 \left[\frac{(4.7 \pm 0.6) + (5.1 \pm 1.6)}{8.3 \pm 1.12} = 1.2 \pm 0.3 \right]_{\text{expt. value}} \quad (33)$$

$$\frac{S(\rho^0 K_s) + S(\omega K_s)}{2} = S(\phi K_s) = -\sin 2\beta \quad (34)$$

$$C(\rho^0 K_s) = -C(\omega K_s) \approx 2r'_c \sin \gamma \sin \delta' + O(r'_c r'_{EW}) \quad (35)$$

where

$$r'_c e^{i\delta'} = \frac{C'}{P'} = \frac{|C'|}{|P'|} e^{i\delta'} \quad (36)$$

We obtain some interesting results from Eqs.(9) and (10). First we note that only W -exchange diagram contributes to $\bar{B}_d^0 \rightarrow K^{*+} K^-, K^+ K^{*-}$ decay channels; neglecting this contribution and also neglecting the W -exchange contribution to $\bar{B}_s^0 \rightarrow \rho^+ \pi^-, \rho^- \pi^+$ in Eqs (13) and (14), we get

$$\begin{aligned} \langle K^{*+} K^- | H_W(s) | \bar{B}_s^0 \rangle &= \langle \rho^+ K^- | H_W(s) | \bar{B}_d^0 \rangle \\ &= |V_{ub}| |V_{ud}^*| |T^{+-}| \left[e^{-i\gamma} \bar{\lambda} \frac{f_K}{f_\pi} - \frac{1}{\bar{\lambda}} r_{+-} e^{i\delta_{+-}} \right] \end{aligned} \quad (37)$$

$$\begin{aligned} r_{+-} &= \frac{|P^{+-}|}{|T^{+-}|} \\ \langle K^+ K^{*-} | H_W(s) | \bar{B}_s^0 \rangle &= \langle \pi^+ K^{*-} | H_W(s) | \bar{B}_d^0 \rangle \\ &= |V_{ub}| |V_{ud}^*| |T^{-+}| \left[e^{-i\gamma} \bar{\lambda} \frac{f_{K^*}}{f_\rho} - \frac{1}{\bar{\lambda}} r_{-+} e^{i\delta_{-+}} \right] \end{aligned} \quad (38)$$

Before, we examine the consequences of $B \rightarrow PV$ decays for various channels obtained above, we discuss the various decay channels of $B \rightarrow P_1 P_2$. For these decays, replace K^* with K in Eqs.(7), (10) and (12). Hence for the decay amplitudes $\bar{A}(\bar{B}_d^0 \rightarrow K^0 \bar{K}^0)$, $A(\bar{B}_s^0 \rightarrow K^0 \bar{K}^0)$; replace P_C by p'_c ; r, δ by r_n, δ'_n in Eqs. (17) and (18) and P_c^{0-}, r_{0-}' and δ_{0-}' by p'_c, r'_c and δ'_c in Eq.(21) for the amplitude $\bar{A}(B^- \rightarrow K^0 K^-)$. The amplitudes for the decays $\bar{B}_s^0 \rightarrow K^+ K^-, B^- \rightarrow \eta_s K^-$ and $B^- \rightarrow \pi^- \bar{K}^0$ are given below.

$$\begin{aligned}
\bar{A}(\bar{B}_s^0 \rightarrow K^+ K^-) &= \bar{A}(\bar{B}_d^0 \rightarrow \pi^+ K^-) \\
&= |V_{cb}| |V_{us}| |t'| e^{-i\gamma} + |V_{cb}| |V_{us}| |p'_c| \\
&= |V_{cb}| |V_{us}| |p'_c| \left[1 + r' e^{i(-\gamma+\delta')} \right]
\end{aligned} \tag{39}$$

$$\bar{A}(B^- \rightarrow K^0 K^-) = |V_{cb}| |V_{cs}| |p'_c| \left[-\bar{\lambda} + r'_c e^{i(-\gamma+\delta'_c)} \right] \tag{40}$$

$$\bar{A}(B^- \rightarrow \eta_s K^-) = |V_{cb}| |V_{cs}| \left[p'_c + \frac{1}{3} p'_{EW} \right]$$

In addition we have

$$\bar{A}(B^- \rightarrow \pi^- \bar{K}^0) \approx |V_{cb}| |V_{cs}| |p'_c| \tag{41}$$

For the observables for the decay $B \rightarrow P_1 P_2$:

$$\langle \Gamma_n \rangle, \Delta_n = \frac{\Gamma(B_d^0 \rightarrow K^0 \bar{K}^0) \pm \Gamma(\bar{B}_d^0 \rightarrow K^0 \bar{K}^0)}{2} \tag{42}$$

$$\langle \Gamma_c \rangle, \Delta_c = \frac{\Gamma(B^+ \rightarrow K^+ \bar{K}^0) \pm \Gamma(B^- \rightarrow K^- K^0)}{2} \tag{43}$$

$$A_{CP}^n = \frac{\Delta_n}{\langle \Gamma_n \rangle} \tag{44}$$

$$A_{CP}^c = \frac{\Delta_c}{\langle \Gamma_c \rangle} \tag{45}$$

we obtain the following results

$$\Gamma(\bar{B}_s^0 \rightarrow K^0 \bar{K}^0) \approx \Gamma(B^- \rightarrow \pi^- \bar{K}^0) \tag{46}$$

$$\frac{\langle \Gamma_n \rangle}{\bar{\lambda}^2 \Gamma(B^- \rightarrow \pi^- \bar{K}^0)} \approx \left[1 - \frac{2r'_n}{\bar{\lambda}} \cos \gamma \cos \delta'_n + \frac{r_n'^2}{\bar{\lambda}^2} \right] \tag{47}$$

$$\frac{\langle \Gamma_c \rangle}{\bar{\lambda}^2 \Gamma(B^- \rightarrow \pi^- \bar{K}^0)} \approx \left[1 - \frac{2r'_c}{\bar{\lambda}} \cos \gamma \cos \delta'_n + \frac{r_c'^2}{\bar{\lambda}^2} \right] \tag{48}$$

$$A_{CP}^{n,c} \approx \frac{\bar{\lambda}^2 \Gamma(B^- \rightarrow \pi^- \bar{K}^0)}{\langle \Gamma_{n,c} \rangle} \left[2 \frac{r'_{n,c}}{\bar{\lambda}} \sin \delta'_{n,c} \sin \gamma \right] \tag{49}$$

Using the following values for the branching ratios: [8]

$$\mathcal{B}(B^- \rightarrow \pi^- \bar{K}^0) = (24.1 \pm 1.7) \times 10^{-6} \quad (50)$$

$$\mathcal{B}(B^- \rightarrow K^0 K^-) = (1.2 \pm 0.32) \times 10^{-6} \quad (51)$$

$$\mathcal{B}(B^0 \rightarrow K^0 \bar{K}^0) = (1.13 \pm 0.38) \times 10^{-6}$$

we get

$$\frac{\langle \Gamma_n \rangle}{\bar{\lambda}^2 \Gamma(B^- \rightarrow \pi^- \bar{K}^0)} = 0.94 \pm 0.32 \quad (52)$$

$$\frac{\langle \Gamma_c \rangle}{\bar{\lambda}^2 \Gamma(B^- \rightarrow \pi^- \bar{K}^0)} = 1.00 \pm 0.25 \quad (53)$$

Hence

$$\left(1 - \frac{2r'_n}{\bar{\lambda}} \cos \gamma \cos \delta'_n + \frac{r'^2_n}{\bar{\lambda}^2}\right) = 0.94 \pm 0.32 \quad (54)$$

$$\left(1 - \frac{2r'_c}{\bar{\lambda}} \cos \gamma \cos \delta'_c + \frac{r'^2_c}{\bar{\lambda}^2}\right) = 1.00 \pm 0.25 \quad (55)$$

In this approximation

$$A_{CP}^{n,c} = \frac{2r'_{n,c}}{\bar{\lambda}} \sin \delta'_{n,c} \sin \gamma \quad (56)$$

With improvement in experimental accuracy, these predictions can be tested

From Eq.(41), we get the results

$$\Gamma(\bar{B}_s^0 \rightarrow K^+ K^-) \approx \Gamma(\bar{B}_d^0 \rightarrow \pi^+ K^-) \quad (57)$$

$$A_{CP}(K^+ K^-) \approx A_{CP}(\pi^+ K^-) = \frac{-2r' \sin \delta'_c \sin \gamma}{1 + 2r' \cos \delta'_c \cos \gamma + r'^2} \quad (58)$$

The experimental values [8] for the decay rates and A_{CP} are

$$\begin{aligned} \Gamma(\bar{B}_d^0 \rightarrow \pi^+ K^-) &= (1.82 \pm 0.08) \times 10^{-5} \\ \Gamma(\bar{B}_s^0 \rightarrow K^+ K^-) &< 5.9 \times 10^{-5} \\ A_{CP}(\pi^+ K^-) &= -0.113 \pm 0.020 \end{aligned}$$

Eqs.(57) and (58) can be tested when the experimental data on $\bar{B}_s^0 \rightarrow K^+ K^-$ will be available.

In order to discuss the mixing induced CP -asymmetries we first give a general expression for the time dependent decay rates for $B^0 \rightarrow f, \bar{f}$ in terms of these asymmetries

$$\Gamma_{\bar{f},f}(t) = e^{-\Gamma t} \frac{1}{2} (R_f + R_{\bar{f}}) (1 \pm A_{CP}) \left\{ \begin{array}{l} 1 + \cos \Delta m t (C_f \pm \Delta C_f) \\ - \sin \Delta m t (S_f \pm \Delta S_f) \end{array} \right\} \quad (59)$$

The decay rates $\bar{\Gamma}_{\bar{f},f}$ can be obtained from Eq.(59) by changing $\cos \Delta m t \rightarrow -\cos \Delta m t$, $\sin \Delta m t \rightarrow -\sin \Delta m t$.

The direct CP asymmetries $C_f \pm \Delta C_f$ and mixing induced CP -asymmetries $S_f \pm \Delta S_f$ in terms of scattering amplitudes are given by

$$\begin{aligned} C_f \pm \Delta C_f &= \frac{|A_{\bar{f},f}|^2 - |\bar{A}_{\bar{f},f}|^2}{|A_{\bar{f},f}|^2 + |\bar{A}_{\bar{f},f}|^2} \\ &= \frac{\pm (R_f - R_{\bar{f}}) + (R_f a_f + R_{\bar{f}} a_{\bar{f}})}{(R_f + R_{\bar{f}}) (1 \pm A_{CP})} \end{aligned} \quad (60)$$

$$(R_f + R_{\bar{f}}) (1 \pm A_{CP}) (S_f \pm \Delta S_f) = 2 \text{Im} \left[e^{-2i\phi_M} A_{\bar{f},f}^* \bar{A}_{\bar{f},f} \right] \quad (61)$$

$$R_f = \frac{|A_{\bar{f}}|^2 + |\bar{A}_f|^2}{2}, \quad R_{\bar{f}} = \frac{|A_f|^2 + |\bar{A}_{\bar{f}}|^2}{2} \quad (62)$$

$$\begin{aligned} R_f a_f &= \frac{1}{2} \left[|A_{\bar{f}}|^2 - |\bar{A}_f|^2 \right], \\ R_{\bar{f}} a_{\bar{f}} &= \frac{1}{2} \left[|A_f|^2 - |\bar{A}_{\bar{f}}|^2 \right] \end{aligned} \quad (63)$$

$$A_{CP} = \frac{R_f a_f - R_{\bar{f}} a_{\bar{f}}}{R_f + R_{\bar{f}}} \quad (64)$$

$$A_{CP}^{\bar{f}} = -a_f, \quad A_{CP}^f = -a_{\bar{f}} \quad (65)$$

Note the weak phase $\phi_M = \beta$ for B_d^0 and $\phi_M = 0$ for B_s^0 . For the case $|\bar{f}\rangle = CP|f\rangle$, we have $R_f = R_{\bar{f}}$, $a_{\bar{f}} = a_f$, $A_{CP} = 0$, $\Delta C_f = 0$, $\Delta S_f = 0$

For this case, we have

$$\frac{\Gamma_f(t) - \bar{\Gamma}_f(t)}{\Gamma_f(t) + \bar{\Gamma}_f(t)} = \cos \Delta m t C_f - \sin \Delta m t S_f \quad (66)$$

$$C_f = \frac{|A_f|^2 - |\bar{A}_f|^2}{|A_f|^2 + |\bar{A}_f|^2} \quad (67)$$

$$S_f = \frac{2 \operatorname{Im} [e^{-2i\phi_M} A_f^* \bar{A}_f]}{|A_f|^2 + |\bar{A}_f|^2} \quad (68)$$

The mixing-induced CP asymmetry $S(\pi^+ K^-)$ is zero, since the decay $\bar{B}_d^0 \rightarrow \pi^- K^+$ is not allowed in the standard Model. However, for the decay $\bar{B}_d^0 \rightarrow K^0 \bar{K}^0$ and $\bar{B}_s^0 \rightarrow K^+ K^-$, the CP asymmetries, can be easily calculated using Eqs. (66,67,68). We get

$$C_n = A_{CP}^n \quad (69)$$

$$S_n = -\frac{\bar{\lambda}^2 \Gamma(B^- \rightarrow \pi^- \bar{K}^0)}{\langle \Gamma_n \rangle} \left\{ \begin{array}{l} \sin 2\beta - 2\frac{r'}{\bar{\lambda}} \sin(\gamma + 2\beta) \cos \delta'_n \\ + \frac{r'^2}{\bar{\lambda}^2} \sin(2\beta + 2\gamma) \end{array} \right\} \quad (70)$$

$$\begin{aligned} S_n(K^+ K^-) &= \frac{-\sin 2\gamma - 2r' \cos \delta'_c \sin \gamma}{1 + 2r' \cos \delta'_c \cos \gamma + r'^2} \\ &\approx -\sin 2\gamma + 2r' \cos \delta'_c \sin \gamma \cos 2\gamma \end{aligned} \quad (71)$$

The mixing induced CP -asymmetry $S(K^+ K^-)$ and direct CP -asymmetry $A_{CP}(\pi^+ K^-)$ would give an alternate way of determining phase γ , when experimental data on $\bar{B}_s^0 \rightarrow K^+ K^-$ will become available.

For the decay $B^0 \rightarrow PV$, we get similar results as for $B^0 \rightarrow P_1 P_2$. In particular, the following results are of particular interest. From Eqs. (29) to (32), we obtain

$$\frac{\langle \Gamma^{0-} \rangle}{\bar{\lambda}^2 \Gamma(B^- \rightarrow \bar{K}^{*0} \pi^-)} = \left[1 - 2\frac{r'_{0-}}{\bar{\lambda}} \cos \gamma \cos \delta'_{0-} + \frac{r_{0-}^2}{\bar{\lambda}^2} \right] \quad (72)$$

$$\frac{\langle \Gamma^{-0} \rangle}{\bar{\lambda}^2 \Gamma(B^- \rightarrow \rho^- \bar{K}^0)} = \left[1 - 2\frac{r'_{-0}}{\bar{\lambda}} \cos \gamma \cos \delta'_{-0} + \frac{r_{-0}^2}{\bar{\lambda}^2} \right] \quad (73)$$

$$a_{CP}^{0-} = \frac{\bar{\lambda}^2 \Gamma(B^- \rightarrow \bar{K}^{*0} \pi^-)}{\langle \Gamma^{0-} \rangle} \left[2\frac{r'_{0-}}{\bar{\lambda}} \sin \gamma \sin \delta'_{0-} \right] \quad (74)$$

$$a_{CP}^{-0} = \frac{\bar{\lambda}^2 \Gamma(B^- \rightarrow \rho^- \bar{K}^0)}{\langle \Gamma^{-0} \rangle} \left[2\frac{r'_{-0}}{\bar{\lambda}} \sin \gamma \sin \delta'_{-0} \right] \quad (75)$$

There is no experimental data on $B^- \rightarrow K^{*0} K^-$ and $B^- \rightarrow K^{*-} \bar{K}^0$ decays to draw conclusion regarding weak phase γ and strong phases δ'_{0-} and δ'_{-0} . However from Eqs.(37) and (38), we obtain observables for which experimental data is available. In particular, we get

$$\Gamma(\bar{B}_s^0 \rightarrow K^{*-} K^+) = \Gamma(\bar{B}^0 \rightarrow K^{*-} \pi^+) \quad (76)$$

$$\Gamma(\bar{B}_s^0 \rightarrow K^{*+} K^-) = \Gamma(\bar{B}^0 \rightarrow \rho^+ K^-) \quad (77)$$

$$A_{CP}^{\pm} = -a'(K^{*-} K^+) = -a'(K^{*-} \pi^+) = \frac{|T^{-+}|^2}{R'_{-+}} \left[2r_{-+} \frac{f_{K^*}}{f_\rho} \sin \gamma \sin \delta_{-+} \right] \quad (78)$$

$$A_{CP}^{\mp} = -a(K^{*+} K^-) = -a'(\rho^+ K^-) = \frac{|T^{+-}|^2}{R'_{+-}} \left[2r_{+-} \frac{f_K}{f_\pi} \sin \gamma \sin \delta_{+-} \right] \quad (79)$$

$$A_{CP}^{\pm} = -a_{-+} = \frac{|T^{-+}|^2}{R_{-+}} [-2r_{-+} \sin \gamma \sin \delta_{-+}] \quad (80)$$

$$A_{CP}^{\mp} = -a_{+-} = \frac{|T^{+-}|^2}{R_{+-}} [-2r_{+-} \sin \gamma \sin \delta_{+-}] \quad (81)$$

where

$$\frac{R'_{-+}}{|T^{-+}|^2} = B'_{-+} = \left[\left(\bar{\lambda} \frac{f_{K^*}}{f_\rho} \right)^2 - 2r_{-+} \frac{f_{K^*}}{f_\rho} \cos \gamma \cos \delta_{-+} + \frac{1}{\bar{\lambda}^2} r_{-+}^2 \right] \quad (82)$$

$$\frac{R'_{+-}}{|T^{+-}|^2} = B'_{+-} = \left[\left(\bar{\lambda} \frac{f_K}{f_\pi} \right)^2 - 2r_{+-} \frac{f_K}{f_\pi} \cos \gamma \cos \delta_{+-} + \frac{1}{\bar{\lambda}^2} r_{+-}^2 \right] \quad (83)$$

$$\frac{R_{-+}}{|T^{-+}|^2} = B_{-+} = [1 + 2r_{-+} \cos \gamma \cos \delta_{-+} + r_{-+}^2] \quad (84)$$

$$\frac{R_{+-}}{|T^{+-}|^2} = B_{+-} = [1 + 2r_{+-} \cos \gamma \cos \delta_{+-} + r_{+-}^2] \quad (85)$$

From above equations, we obtain the results

$$\frac{a'_{-+}}{a_{-+}} = - \left(\frac{f_{K^*}}{f_\rho} \right) \left(\frac{R_{-+}}{R'_{-+}} \right) \quad (86)$$

$$\frac{a'_{+-}}{a_{+-}} = - \left(\frac{f_K}{f_\pi} \right) \left(\frac{R_{+-}}{R'_{+-}} \right) \quad (87)$$

$$r_{-+}^2 = \bar{\lambda}^2 \left[B_{-+} \frac{R'_{-+}}{R_{-+}} + (B_{-+} - 1) \frac{f_{K^*}}{f_\rho} \right] \quad (88)$$

$$r_{+-}^2 = \bar{\lambda}^2 \left[B_{+-} \frac{R'_{+-}}{R_{+-}} + (B_{+-} - 1) \frac{f_K}{f_\pi} \right] \quad (89)$$

In the end, we discuss the relations between various observables defined in Eqs.(59-66) for the decays $\bar{B}_d^0 \rightarrow \rho^- \pi^+, \rho^+ \pi^-$ and $\bar{B}_s^0 \rightarrow K^{*-} K^+, K^{*+} K^-$. The CP violating asymmetry A_{CP} , the direct CP violation C , the dilution parameter ΔC , the mixing induced CP violation S and dilution parameter ΔS for the $\rho\pi$ and K^*K decay channels, are related to each other by $SU(3)$. We obtain the following results for the various observables, using Eqs. (37) and (38); dashed quantites refer to $\bar{B}_s^0 \rightarrow K^{*-} K^+, K^{*+} K^-$ and undashed to $\bar{B}_d^0 \rightarrow \rho^- \pi^+, \rho^+ \pi^-$

$$A'_{CP} = -\frac{1}{2} \frac{R_{-+} + R_{+-}}{R'_{-+} + R'_{+-}} \left[\left(\frac{f_{K^*}}{f_\rho} + \frac{f_K}{f_\pi} \right) A_{CP} + \left(\frac{f_{K^*}}{f_\rho} - \frac{f_K}{f_\pi} \right) (C + A_{CP} \Delta C) \right] \quad (90)$$

$$C' + A'_{CP} \Delta C' = -\frac{1}{2} \frac{R_{-+} + R_{+-}}{R'_{-+} + R'_{+-}} \left[\left(\frac{f_{K^*}}{f_\rho} + \frac{f_K}{f_\pi} \right) (C + A_{CP} \Delta C) + \left(\frac{f_{K^*}}{f_\rho} - \frac{f_K}{f_\pi} \right) A_{CP} \right] \quad (91)$$

$$\Delta C + A_{CP} C = \frac{R_{-+} - R_{+-}}{R_{-+} + R_{+-}}; \Delta C' + A'_{CP} C' = \frac{R'_{-+} - R'_{+-}}{R'_{-+} + R'_{+-}} \quad (92)$$

Taking $\delta_t = (\delta_T^{+-} - \delta_T^{-+}) = 0$ [9], we get for mixing induced CP - violating parameters S and ΔS , the following results: (neglecting terms of the order $r_{-+} r_{+-}$)

$$\frac{1}{t} (B_{-+} + t^2 B_{+-}) [\Delta S' + A'_{CP} S'] \approx \frac{R_{-+} + R_{+-}}{R'_{-+} + R'_{+-}} 2 \cos \gamma \left[-\frac{f_{K^*}}{f_\rho} r_{+-} \sin \delta_{+-} + \frac{f_K}{f_\pi} r_{-+} \cos \delta_{-+} \right] \quad (93)$$

$$\frac{1}{t} (B_{-+} + t^2 B_{+-}) [S + A_{CP} \Delta S] \approx 2 \left[\frac{-\sin(2\beta + 2\gamma) - \sin(2\beta + \gamma) \times}{(r_{-+} \cos \delta_{-+} + r_{+-} \cos \delta_{+-})} \right] \quad (94)$$

$$\frac{1}{t} (B_{-+} + t^2 B_{+-}) [\Delta S + A_{CP} S] \approx 2 \cos(2\beta + \gamma) [r_{+-} \sin \delta_{+-} - r_{-+} \sin \delta_{-+}] \quad (95)$$

$$\frac{1}{t} (B_{-+} + t^2 B_{+-}) [S' + A'_{CP} \Delta S'] \approx \frac{R_{-+} + R_{+-}}{R'_{-+} + R'_{+-}} 2 \left[\frac{-\bar{\lambda}^2 f_{K^*} f_K \sin 2\gamma}{f_\rho f_\pi} + \sin \gamma \left(\frac{\frac{f_K}{f_\pi} r_{-+} \cos \delta_{-+}}{+\frac{f_{K^*}}{f_\rho} r_{+-} \cos \delta_{+-}} \right) \right] \quad (96)$$

In order to make contact with the experimental data, we use the data of reference [8]. Using the following experimental values

$$R_{-+} + R_{+-} = (22.8 \pm 2.5) \times 10^{-6} \\ R'_{-+} = (11.8 \pm 1.5) \times 10^{-6}, R'_{+-} = (8.5 \pm 2.8) \times 10^{-6} \quad (97)$$

$$a_{-+} = 0.15 \pm 0.08, a_{+-} = 0.53 \pm 0.13 \\ C = 0.30 \pm 0.13, \Delta C = 0.33 \pm 0.13 \quad (98)$$

we obtain

$$A_{CP} = -0.087 \pm 0.07, R_{-+} = (14.9 \pm 2.2) \times 10^{-6}, R_{+-} = (7.9 \pm 1.8) \times 10^{-6} \quad (99)$$

Hence using the above values we get from Eqs.(86), (87)

$$a'_{-+} = -0.20 \pm 0.11 (-0.05 \pm 0.14), a'_{+-} = -0.60 \pm 0.41 (-0.26 \pm 0.15)$$

where the numbers within the brackets are experimental values.

In order to obtain the values of r_{-+} and r_{+-} from Eqs.(88) and (89) we need the values for B_{-+} and B_{+-} . Using [9],

$$B_{-+} = 1.03 \pm 0.12, B_{+-} = 1.05 \pm 0.12 \quad (100)$$

we obtain

$$r_{-+}^2 = 0.044 \pm 0.014; r_{-+} = 0.21 \pm 0.04; 0.17 \leq r_{-+} \leq 0.24 \quad (101)$$

$$r_{+-}^2 = 0.063 \pm 0.038, r_{+-} = 0.25 \pm 0.06; 0.19 \leq r_{+-} \leq 0.30 \quad (102)$$

to be compared with the values given in reference [7]. The following remarks are in order. The values given in Eq.(100) has been obtained from $B_{-+} = \frac{R_{-+}}{|T_{-+}|^2}$, $B_{+-} = \frac{R_{+-}}{|T_{+-}|^2}$, using factorization for the tree amplitude T_{-+} with $|V_{ub}| = 3.45 \times 10^{-3}$, $f_+(m_\rho^2) = 0.26$ and $t^2 = 0.52$

Using the above values of r_{-+} , r_{+-} , B_{-+} and B_{+-} , we get from Eqs.(80,81) and (84,85)

$$x_{-+} \equiv \sin \gamma \sin \delta_{-+} = 0.37 \pm 0.21, \quad 0.16 \leq x_{-+} \leq 0.58 \quad (103)$$

$$x_{+-} \equiv \sin \gamma \sin \delta_{+-} = 1.11 \pm 0.68, \quad 0.43 \leq x_{+-} \leq 1 \quad (104)$$

$$z_{-+} \equiv \cos \gamma \cos \delta_{-+} = -0.03 \pm 0.31, \quad -0.34 \leq z_{-+} < 0.28 \quad (105)$$

$$z_{+-} \equiv \cos \gamma \cos \delta_{+-} = -0.03 \pm 0.30, \quad -0.33 \leq z_{+-} < 0.27 \quad (106)$$

From Eqs.(103) and (104), we obtain the following limits on the final state strong phases, for γ in the range $51^\circ \leq \gamma \leq 75^\circ$ [8]

$$10^\circ \leq \delta_{-+} \leq 48^\circ \text{ or } 170^\circ \geq \delta_{-+} \geq 132^\circ$$

$$27^\circ \leq \delta_{+-} \leq 90^\circ \text{ or } 153^\circ \geq \delta_{+-} \geq 90^\circ$$

We note that x and z satisfy the following equations

$$\frac{x^2}{\sin^2 \gamma} + \frac{z^2}{\cos^2 \gamma} = 1 \quad (107)$$

$$\frac{x^2}{\sin^2 \delta} + \frac{z^2}{\cos^2 \delta} = 1$$

Figs(1,2) display these ellipses in (x, z) plane for x_{-+}, z_{-+} and x_{+-}, z_{+-} respectively for various values of γ and δ . The experimental data restrict the allowed area in (x, z) plane to the dark shaded regions in Figs(1,2). It is clear from Fig 1, that for $z_{-+} < 0$, $\gamma > 65^\circ$, whereas $\gamma > 70^\circ$ for $z_{-+} > 0$. From Figs 1 and 2, we also note that $\delta_{-+} \leq 40^\circ$ for $z_{-+} > 0$ or $\delta_{-+} \geq 140^\circ$ for $z_{-+} < 0$, whereas $\delta_{+-} \geq 25^\circ$ or $\delta_{+-} \leq 155^\circ$.

Using the experimental values given in Eqs.(97) and (98), we obtain

$$\frac{R_{-+} + R_{+-}}{R'_{-+} + R'_{+-}} = 1.12 \pm 0.22 \quad (108)$$

$$A'_{CP} = 0.14 \pm 0.09 \quad (109)$$

$$C' + A'_{CP} \Delta C' = -0.34 \pm 0.17 \quad (110)$$

Since the experimental values for mixing induced asymmetries S and ΔS have large errors, there is no point in giving the numerical values for S' and $\Delta S'$.

To conclude:

Using rotation in $SU(3)$ space, a set of relations between various decay modes of B mesons have been derived. In this way, we have avoided use of group representation of $SU(3)$ symmetry. In particular we have related the decay modes of $\bar{B}_d^0$ and $\bar{B}_s^0$ to $K\bar{K}, K^*\bar{K}, \bar{K}^*K$ and $\bar{B}_s^0 \rightarrow K^+K^-$ to $\bar{B}_d^0 \rightarrow \pi^+K^-$. Due to lack of experimental data, some of the relations cannot be tested. However for $B^0 \rightarrow \rho K, \omega K, \phi K$ decays we get

$$\frac{\langle \Gamma \rangle_{\omega K} + \langle \Gamma \rangle_{\rho^0 K}}{\langle \Gamma \rangle_{\phi K}} \approx 1, \quad \frac{S(\rho^0 K_s) + S(\omega K_s)}{2} = S(\phi K_s) = -\sin 2\beta \quad (111)$$

$$C(\rho^0 K_s) = -C(\omega K_s) \quad (112)$$

$$f_\rho - \frac{1}{3}f_\omega - \frac{2}{3}f_\phi = 0 \quad (113)$$

In more details, we have expressed the decay amplitudes $\langle K^{*-}K^+ | H_w(s) | \bar{B}_s^0 \rangle = \langle K^{*-}\pi^+ | H_w(s) | \bar{B}_d^0 \rangle$ and $\langle K^{*+}K^- | H_w(s) | \bar{B}_s^0 \rangle = \langle \rho^+K^- | H_w(s) | \bar{B}_d^0 \rangle$ in terms of the decay parameters of $\bar{B}_d^0 \rightarrow \rho^-\pi^+$ and $\bar{B}_d^0 \rightarrow \rho^+\pi^-$ respectively. In particular in Eqs.(88) and (89), we have obtained $r_{-+}(r_{+-})$ in terms of two unknown parameters $B_{-+}(B_{+-})$ as all other terms in these equations are known experimentally. These equations follow from broken $SU(3)$. $SU(3)$ is a good symmetry except for the masses and the matrix elements of weak currents between vacuum and pseudoscalar or vector mesons belonging to octet or nonet representations of $SU(3)$. Thus $SU(3)$ breaking effects are taken care of by using physical values for masses and for f_π, f_K, f_ρ and f_{K^*} . Parameters $B_{-+}(B_{+-})$ are determined using factorization for tree graph. Having determined $r_{-+}(r_{+-})$, we obtain $z_{-+}(z_{+-})$ and $x_{-+}(x_{+-})$ from Eqs.(84), (85), (80) and (81). From x_{-+} and x_{+-} following bounds on strong phases are obtained

$$10^\circ \leq \delta_{-+} \leq 48^\circ, \quad 27^\circ \leq \delta_{+-} \leq 90^\circ \quad (114)$$

For $z_{-+}, z_{+-} < 0$, the angles lie in the second quadrant. Further from Fig(1), we get lower limit on γ to be 65° , and for the final state phases we get

$$\delta_{-+} < 40^\circ (\delta_{-+} \geq 160^\circ)$$

Figure 1: Plot of x_{-+} versus z_{-+} for various values of γ (solid curves) and δ (dotted curves). The dark shaded area is the region allowed by experimental data.

Figure 2: Plot of x_{+-} versus z_{+-} for various values of γ (solid curves) and δ (dotted curves). The dark shaded area is the region allowed by experimental data.

From Fig 2, we get

$$\delta_{+-} > 25^\circ (\delta_{+-} < 155^\circ)$$

Acknowledgments:

The author acknowledges a research grant provided by the Higher Education Commission of Pakistan to him as a Distinguished National Professor. The author also wishes to thank Mr. Ijaz Ahmed for drawing the figures.

References

- [1] For recent reviews, see for instance, M.Gronau, arXiv:hep-ph/0510153 v3, 2005; M.Gronau, E. Lunghi and D.Wyler, arXiv:hep-ph/0410170 v3, 2004; C. W. Baur et al. arXiv: hep-ph/0401188 v2, 2004; M. Beneke and M. Neubert, Nucl. Phys. B 675 (2003) 33; U. Nierste, arXiv:hep-ph/0511125 v1, 2005; F.Forti, arXiv:hep-ex/0511010 v1, 2005; S. Safir, Proc. Sci, HEP2005: 246 (2005), hep-ph/0512014, M.Gronau and Jonathan.L.Rosner, hep-ph/0307095, M.Gronau and Jonathan.L.Rosner, hep-ph/0610227.
- [2] See for example: Fayyazuddin and Riazuddin, A Modern Introduction to Particle Physics Second Edition Page (134) World Scientific (2000)
- [3] M.Gronau, Phys.Rev.D62 (2000) 014031.
- [4] N.G.Deshpande, Xiao-Gang He and Jian-Qing Shi, hep-ph/0002260 v2, Phys.Rev. D62 (2000) 034018
- [5] T.Das, V.S. Mathur and S.Okubo, Phys.Rev.Lett; 19 (1967) 470. See also J.J. Sakurai, Currents and Mesons, The University of Chicago Press (1969)
- [6] See for example: M.Beneke and M.Nuebert, Nucl.Phys B675,333(2003)
- [7] M.Gronau and J.Zupan, hep-ph/0407002, 2004;
- [8] Particle Data Group, W-M.Yao *etal.* Journal of Physics G **33**,1 (2006)
- [9] Fayyazuddin hep-ph/0604008 v3, 2007.

Figure Captions:

Figure1: Plot of x_{-+} versus z_{-+} for various values of γ (solid curves) and δ (dotted curves). The dark shaded area is the region allowed by experimental data.

Figure2: Plot of x_{+-} versus z_{+-} for various values of γ (solid curves) and δ (dotted curves). The dark shaded area is the region allowed by experimental data.

This figure "Figure1.JPG" is available in "JPG" format from:

<http://arXiv.org/ps/hep-ph/0607310v2>

This figure "Figure2.JPG" is available in "JPG" format from:

<http://arXiv.org/ps/hep-ph/0607310v2>